

# Crepant Resolutions of Gorenstein Toric Singularities and Upper Bound Theorem

\*Dimitrios I. Dais

Mathematics Department, Section of Algebra and Geometry, University of Ioannina

GR-45110 Ioannina, Greece. E-mail: ddais@cc.uoi.gr

## Abstract

A necessary condition for the existence of torus-equivariant crepant resolutions of Gorenstein toric singularities can be derived by making use of a variant of the classical Upper Bound Theorem which is valid for simplicial balls.

## 1 Introduction

Let  $d$  be a positive integer,  $\sigma \subset \mathbb{R}^{d+1}$  a rational,  $(d+1)$ -dimensional strongly convex polyhedral cone (w.r.t. the lattice  $\mathbb{Z}^{d+1}$ ), and

$$U_\sigma = \text{Spec} \left( \mathbb{C} [\sigma^\vee \cap (\mathbb{Z}^{d+1})^\vee] \right)$$

the associated affine toric variety, where  $\sigma^\vee$  denotes the dual of  $\sigma$ . (For the usual notions of toric geometry, see [7]). As it is known (see e.g. [10, §6]):

**Theorem 1.1**  *$U_\sigma$  is Gorenstein if and only if the set  $\text{Gen}(\sigma)$  of the minimal generating integral vectors of  $\sigma$  lies on a “primitive” affine hyperplane, i.e., iff*

$$\text{Gen}(\sigma) \subset \mathbf{H}_\sigma = \{ \mathbf{x} \in \mathbb{R}^{d+1} \mid \langle \mathbf{m}_\sigma, \mathbf{x} \rangle = 1 \},$$

where  $\mathbf{m}_\sigma \in (\mathbb{Z}^{d+1})^\vee$  is a primitive vector belonging to the dual lattice.

**Remark 1.2** (i) In this case,  $\sigma$  supports the  $d$ -dimensional lattice polytope

$$P_\sigma = \{ \mathbf{x} \in \sigma \mid \langle \mathbf{m}_\sigma, \mathbf{x} \rangle = 1 \} \subset \mathbf{H}_\sigma \cong \mathbb{R}^d \quad (1.1)$$

(w.r.t. the lattice  $\mathbf{H}_\sigma \cap \mathbb{Z}^{d+1} \cong \mathbb{Z}^d$ ).

(ii) In fact, every lattice  $d$ -polytope  $P \subset \mathbb{R}^d$  can be considered as supported by a cone

$$\sigma_P = \{ (r, r\mathbf{x}) \in \mathbb{R} \oplus \mathbb{R}^d \mid \mathbf{x} \in P, r \in \mathbb{R}_{\geq 0} \} \subset \mathbb{R}^{d+1}$$

so that  $U_{\sigma_P}$  is Gorenstein.

---

\*GSRT-fellow supported by the European Union and the Greek Ministry of Research and Technology.

2000 *Mathematics Subject Classification*. Primary 14M25, 52B20; Secondary 14B05, 52B05, 52B11.

The  $(d+1)$ -dimensional Gorenstein toric singularities<sup>1</sup>  $(U_\sigma, \text{orb}(\sigma))$  constructed by cones  $\sigma$  which support lattice  $d$ -polytopes  $P = P_\sigma$  are to be subdivided into three distinct classes<sup>2</sup>:

(A) *Terminal singularities* (whenever  $P$  is an elementary polytope but not a basic simplex).

(B) *Canonical, non-terminal singularities which do not admit any crepant resolution* (i.e., for which  $P$  is a non-elementary polytope having no basic triangulations).

(C) *Canonical, non-terminal singularities admitting crepant resolutions* (i.e., for which  $P$  is a non-elementary polytope possessing *at least one* basic triangulation).

**Comments.** A complete classification of the members of class (A) (up to analytic isomorphism) is obviously equivalent to the classification of elementary polytopes (up to lattice automorphism). For constructions of several families belonging to (C), the reader is referred to [1], [2], [3], [4], [5]. In fact, for one- or two-parameter Gorenstein cyclic quotient singularities, it is possible to decide definitely if they belong to class (A), (B) or (C), by just checking some concrete number-theoretic (necessary and sufficient existence-) conditions (see [3] and [2], respectively). On the other hand, for *general* Gorenstein toric (not necessarily quotient-) singularities, a *necessary condition* for the existence of crepant resolutions can be created via an UBT for simplicial balls, as we shall see below in Thm. 3.1. Hence, its “violation” may be used to produce families of such singularities belonging to (B).

## 2 Basic facts about UBT's

NOTATION. (i) The **f**-vector  $\mathbf{f}(\mathcal{S}) = (f_0(\mathcal{S}), f_1(\mathcal{S}), \dots, f_{d-1}(\mathcal{S}))$  of a polyhedral  $(d-1)$ -complex  $\mathcal{S}$  is defined by setting for all  $i$ ,  $0 \leq i \leq d-1$ ,

$$f_i(\mathcal{S}) := \# \{i\text{-dimensional faces of } \mathcal{S}\}$$

(under the usual conventional extension:  $f_{-1}(\mathcal{S}) := 1$ ). The coordinates of the **h**-vector  $\mathbf{h}(\mathcal{S}) = (h_0(\mathcal{S}), h_1(\mathcal{S}), \dots, h_{d-1}(\mathcal{S}), h_d(\mathcal{S}))$  of such an  $\mathcal{S}$  are defined by the equations

$$h_j(\mathcal{S}) = \sum_{i=0}^j (-1)^{j-i} \binom{d-i}{d-j} f_{i-1}(\mathcal{S}). \quad (2.1)$$

(ii) For a  $d$ -dimensional polytope  $P$ , the *boundary complex*  $\mathcal{S}_{\partial P}$  of  $P$  is defined to be the  $(d-1)$ -dimensional polyhedral complex consisting of the proper faces of  $P$  together with  $\emptyset$  and having  $\partial P$  as its support.  $\mathcal{S}_{\partial P}$  is a polyhedral  $(d-1)$ -sphere.  $\mathcal{S}_{\partial P}$  is a geometric pure simplicial complex (in fact, a simplicial  $(d-1)$ -sphere) if and

---

<sup>1</sup>Without loss of generality, we may henceforth assume that the cones  $\sigma \subset \mathbb{R}^{d+1}$  are  $(d+1)$ -dimensional, and that the singularities under consideration have maximal splitting codimension. (The orbit  $\text{orb}(\sigma) \in U_\sigma$  is the unique fixed closed point under the usual torus-action on  $U_\sigma$ .)

<sup>2</sup>A lattice polytope  $P$  is called *elementary* if the lattice points belonging to it are exactly its vertices. A lattice simplex is said to be *basic* (or *unimodular*) if its vertices constitute a part of a  $\mathbb{Z}$ -basis of the reference lattice (or equivalently, if its relative, normalized volume equals 1). A lattice triangulation  $\mathcal{T}$  of a lattice polytope  $P$  is defined to be *maximal* (resp. *basic*), if it consists only of elementary (resp. basic) simplices.

only if  $P$  is a simplicial polytope. The  $\mathbf{f}$ -vector  $\mathbf{f}(P)$  of a  $d$ -polytope  $P$  is by definition the  $\mathbf{f}$ -vector  $\mathbf{f}(\mathcal{S}_{\partial P})$  of its boundary complex.

(iii) We denote by  $\text{CycP}_d(k)$  the *cyclic  $d$ -polytope* with  $k$  vertices. As it is known, the number of its facets equals

$$f_{d-1}(\text{CycP}_d(k)) = \binom{k - \lceil \frac{d}{2} \rceil}{\lfloor \frac{d}{2} \rfloor} + \binom{k - 1 - \lceil \frac{d-1}{2} \rceil}{\lfloor \frac{d-1}{2} \rfloor} \quad (2.2)$$

This follows from Gale's evenness condition and the fact that  $\text{CycP}_d(k)$  is  $\lfloor \frac{d}{2} \rfloor$ -neighbourly (cf. [13, p. 24]).

(iv) Classical UB and LB-Theorems for simplicial spheres (see [9] and [6]):

**Theorem 2.1 (Stanley's Upper Bound Theorem for Simplicial Spheres)**

*The  $\mathbf{f}$ -vector coordinates of a simplicial  $(d-1)$ -sphere  $\mathcal{S}$  with  $f_0(\mathcal{S}) = k$  vertices satisfy the following inequalities:*

$$f_i(\mathcal{S}) \leq f_i(\text{CycP}_d(k)) \quad , \quad \forall i \quad , \quad 0 \leq i \leq d-1.$$

**Theorem 2.2 (Lower Bound Theorem for Simplicial Spheres)** *The  $\mathbf{h}$ -vector coordinates of a simplicial  $(d-1)$ -sphere  $\mathcal{S}$  with  $f_0(\mathcal{S}) = k$  vertices satisfy the following inequalities:*

$$h_1(P) = k - d \leq h_i(P) \quad , \quad \forall i \quad , \quad 2 \leq i \leq d.$$

Besides them we need certain variants for simplicial *balls*.

**Proposition 2.3 (“ $\mathbf{h}$  of  $\partial$ ”-Lemma)** *Let  $\mathcal{S}$  be a  $d$ -dimensional Cohen-Macaulay closed pseudomanifold with non-empty boundary  $\partial\mathcal{S}$ . Then*

$$h_{i-1}(\partial\mathcal{S}) - h_i(\partial\mathcal{S}) = h_{(d+1)-i}(\mathcal{S}) - h_i(\mathcal{S}) \quad , \quad \forall i \quad , \quad 0 \leq i \leq d+1 \quad (2.3)$$

(under the convention:  $h_{-1}(\partial\mathcal{S}) = 0$ ).

*Proof.* See Stanley ([12, 2.3]). □

Working with *Buchsbaum complexes*, Schenzel [8] proved the following:

**Theorem 2.4 (Schenzel's Upper Bound Theorem)** *Let  $\mathcal{S}$  be a  $d$ -dimensional Buchsbaum complex<sup>3</sup> having  $f_0(\mathcal{S}) = \mathbf{b}$  vertices. Then for all  $i$ ,  $0 \leq i \leq d+1$ , the  $\mathbf{h}$ -vector coordinates of  $\mathcal{S}$  satisfy the inequalities*

$$h_i(\mathcal{S}) \leq \binom{\mathbf{b}-d+i-2}{i} - (-1)^i \binom{d+1}{i} \sum_{j=-1}^{i-2} (-1)^j \dim_{\mathbf{k}} \tilde{H}_j(\mathcal{S}; \mathbf{k}) \quad (2.4)$$

(where  $\tilde{H}_j(\mathcal{S}; \mathbf{k})$  are the reduced homology groups of  $\mathcal{S}$  with coefficients in a field  $\mathbf{k}$ .)

<sup>3</sup>A simplicial complex  $\mathcal{S}$  is a Buchsbaum complex over a field  $\mathbf{k}$  if and only if it is pure and the localizations  $\mathbf{k}[\mathcal{S}]_{\mathfrak{p}}$  of  $\mathbf{k}[\mathcal{S}]$  w.r.t. prime ideals  $\mathfrak{p} \neq \mathbf{k}[\mathcal{S}]_+$  ( $= \bigoplus_{\nu > 0} (\mathbf{k}[\mathcal{S}]_{\nu})$ ) are Cohen-Macaulay. (For instance, homology  $d$ -manifolds without boundary, or homology  $d$ -manifolds whose boundary is a homology  $(d-1)$ -manifold without boundary, are Buchsbaum). Moreover,  $\mathcal{S}$  is Cohen-Macaulay over  $\mathbf{k}$  if and only if  $\mathcal{S}$  is Buchsbaum over  $\mathbf{k}$  and  $\dim_{\mathbf{k}} \tilde{H}_j(\mathcal{S}; \mathbf{k}) = 0$ , for all  $i, 0 \leq i \leq d-1$ , while  $\dim_{\mathbf{k}} \tilde{H}_d(\mathcal{S}; \mathbf{k}) = (-1)^d \tilde{\chi}(\mathcal{S})$ , with  $\tilde{\chi}(\mathcal{S})$  the reduced Euler characteristic.

**Corollary 2.5** *Let  $\mathcal{S}$  denote a simplicial  $d$ -dimensional **ball** with  $f_0(\mathcal{S}) = \mathfrak{b}$  vertices. Then for all  $i$ ,  $0 \leq i \leq d$ , the  $\mathbf{f}$ -vector of  $\mathcal{S}$  satisfies the following inequalities:*

$$f_i(\mathcal{S}) \leq f_i(\text{CycP}_{d+1}(\mathfrak{b})) - \sum_{j=d-i}^{\lfloor \frac{d}{2} \rfloor} \binom{j}{d-i} (\mathfrak{h}_j(\partial\mathcal{S}) - \mathfrak{h}_{j-1}(\partial\mathcal{S})), \quad (2.5)$$

*Proof.* Introduce the auxiliary vector  $\tilde{\mathbf{h}}(\mathcal{S}) = (\tilde{\mathfrak{h}}_0(\mathcal{S}), \dots, \tilde{\mathfrak{h}}_{d+1}(\mathcal{S}))$  with

$$\tilde{\mathfrak{h}}_i(\mathcal{S}) := \begin{cases} \mathfrak{h}_i(\mathcal{S}) & , \text{ for } 0 \leq i \leq \lfloor \frac{d+1}{2} \rfloor \\ \mathfrak{h}_i(\mathcal{S}) - (\mathfrak{h}_{d-i}(\partial\mathcal{S}) - \mathfrak{h}_{d+1-i}(\partial\mathcal{S})) & , \text{ for } \lfloor \frac{d+1}{2} \rfloor + 1 \leq i \leq d+1 \end{cases}$$

Since  $\mathcal{S}$  is Cohen-Macaulay,  $\mathcal{S}$  is a Buchsbaum complex. Moreover, all reduced homology groups  $\tilde{H}_j(\mathcal{S}; \mathbf{k})$  are trivial, which means that

$$\mathfrak{h}_i(\mathcal{S}) \leq \mathfrak{h}_i(\text{CycP}_{d+1}(\mathfrak{b})) = \binom{\mathfrak{b}-d+i-2}{i}, \quad \forall i, \quad 0 \leq i \leq \lfloor \frac{d+1}{2} \rfloor,$$

by (2.4). On the other hand, (2.3) implies for the coordinates of  $\tilde{\mathbf{h}}(\mathcal{S})$ :

$$\tilde{\mathfrak{h}}_i(\mathcal{S}) = \tilde{\mathfrak{h}}_{(d+1)-i}(\mathcal{S}), \quad \forall i, \quad 0 \leq i \leq d+1,$$

and therefore

$$\tilde{\mathfrak{h}}_i(\mathcal{S}) \leq \mathfrak{h}_i(\text{CycP}_{d+1}(\mathfrak{b})), \quad \forall i, \quad 0 \leq i \leq d+1. \quad (2.6)$$

Hence,

$$\begin{aligned} f_i(\mathcal{S}) &= \sum_{j=0}^{i+1} \binom{d+1-j}{d-i} \mathfrak{h}_j(\mathcal{S}) \\ &= \sum_{j=0}^{i+1} \binom{d+1-j}{d-i} \tilde{\mathfrak{h}}_j(\mathcal{S}) + \sum_{j=\lfloor \frac{d}{2} \rfloor + 1}^{i+1} \binom{d+1-j}{d-i} (\mathfrak{h}_{d-j}(\partial\mathcal{S}) - \mathfrak{h}_{d+1-j}(\partial\mathcal{S})) \\ &= \sum_{j=0}^{i+1} \binom{d+1-j}{d-i} \tilde{\mathfrak{h}}_j(\mathcal{S}) + \sum_{j=d-i}^{\lfloor \frac{d}{2} \rfloor} \binom{j}{d-i} (\mathfrak{h}_{j-1}(\partial\mathcal{S}) - \mathfrak{h}_j(\partial\mathcal{S})) \\ &\quad [\text{by interchanging } (d+1)-j \text{ and } j, \text{ and using} \\ &\quad \text{the Dehn-Sommerville relations for } \mathbf{h}(\partial\mathcal{S})] \\ &\leq \sum_{j=0}^{i+1} \binom{d+1-j}{d-i} \mathfrak{h}_i(\text{CycP}_{d+1}(\mathfrak{b})) + \\ &\quad + \sum_{j=d-i}^{\lfloor \frac{d}{2} \rfloor} \binom{j}{d-i} (\mathfrak{h}_{j-1}(\partial\mathcal{S}) - \mathfrak{h}_j(\partial\mathcal{S})) \quad [\text{by (2.6)}] \\ &= f_i(\text{CycP}_{d+1}(\mathfrak{b})) - \sum_{j=d-i}^{\lfloor \frac{d}{2} \rfloor} \binom{j}{d-i} (\mathfrak{h}_j(\partial\mathcal{S}) - \mathfrak{h}_{j-1}(\partial\mathcal{S})) \end{aligned}$$

for all  $i, 0 \leq i \leq d$ .  $\square$

**Corollary 2.6** *Let  $\mathcal{S}$  be a simplicial  $d$ -ball with  $f_0(\mathcal{S}) = \mathfrak{b}$  vertices. Suppose that  $f_0(\partial\mathcal{S}) = \mathfrak{b}'$ . Then:*

$$f_d(\mathcal{S}) \leq f_d(\text{CycP}_{d+1}(\mathfrak{b})) - (\mathfrak{b}' - d) \quad (2.7)$$

*Proof.* For  $i = d$ , (2.5) gives

$$\begin{aligned} f_d(\mathcal{S}) &\leq f_d(\text{CycP}_{d+1}(\mathfrak{b})) - \sum_{j=0}^{\lfloor \frac{d}{2} \rfloor} (\mathfrak{h}_j(\partial\mathcal{S}) - \mathfrak{h}_{j-1}(\partial\mathcal{S})) \\ &= f_d(\text{CycP}_{d+1}(\mathfrak{b})) - \mathfrak{h}_{\lfloor \frac{d}{2} \rfloor}(\partial\mathcal{S}) \\ &\leq f_d(\text{CycP}_{d+1}(\mathfrak{b})) - \mathfrak{h}_1(\partial\mathcal{S}), \end{aligned}$$

where the latter inequality comes from the LBT 2.2 for the simplicial sphere  $\partial\mathcal{S}$ . Now obviously,  $\mathfrak{h}_1(\partial\mathcal{S}) = \mathfrak{b}' - d$ .  $\square$

### 3 Crepant Resolutions and UBT

Let  $(U_\sigma, \text{orb}(\sigma))$  be a Gorenstein toric singularity as in §1 (cf. Thm. 1.1).

**Theorem 3.1 (Necessary Existence Condition)** *If  $U_\sigma$  admits a crepant desingularization, then the normalized volume of the lattice polytope  $P_\sigma$  (defined in (1.1)) has the following upper bound<sup>4</sup>*

$$\text{Vol}_{\text{norm}}(P_\sigma) \leq f_d(\text{CycP}_{d+1}(\#(P_\sigma \cap \mathbb{Z}^d))) - (\#(\partial P_\sigma \cap \mathbb{Z}^d) - d) \quad (3.1)$$

*Proof.* If  $U_\sigma$  admits a crepant desingularization, then there must be a basic triangulation, say  $\mathcal{T}$  of  $P_\sigma$ . Since this  $\mathcal{T}$  is, in particular, maximal, we have

$$\text{vert}(\mathcal{T}) = P_\sigma \cap \mathbb{Z}^d, \quad \text{vert}(\partial\mathcal{T}) = \partial P_\sigma \cap \mathbb{Z}^d. \quad (3.2)$$

On the other hand,

$$\text{Vol}_{\text{norm}}(P_\sigma) = f_d(\mathcal{T}). \quad (3.3)$$

Finally, since  $\mathcal{T}$  is a simplicial  $d$ -ball, one deduces (3.1) from (2.7), (3.2), (3.3).  $\square$

**Example 3.2** Let

$$\sigma = \mathbb{R}_{\geq 0} e_1 + \mathbb{R}_{\geq 0} e_2 + \mathbb{R}_{\geq 0} e_3 + \mathbb{R}_{\geq 0} (-3, -7, -9, 20) \subset \mathbb{R}^4$$

be the four-dimensional cone supporting the lattice 3-simplex

$$\mathbf{s}_\sigma = \text{conv}(\{e_1, e_2, e_3, (-3, -7, -9, 20)\}) = \{\mathbf{x} \in \sigma \mid \langle \mathbf{m}_\sigma, \mathbf{x} \rangle = 1\},$$

---

<sup>4</sup>By abuse of notation, we write  $\mathbb{Z}^d$  instead of  $\mathbf{H}_\sigma \cap \mathbb{Z}^{d+1}$  ( $\cong \mathbb{Z}^d$ )

where  $\mathbf{m}_\sigma = (1, 1, 1, 1)$ . Obviously,  $\text{Vol}_{\text{norm}}(\mathbf{s}_\sigma) = 20$ . On the other hand, since  $\mathbf{s}_\sigma$  has 8 lattice points (its 4 vertices and further 4 integer points lying in its relative interior), computing the right-hand side of (3.1), we obtain by (2.2):

$$\binom{8 - \lceil \frac{4}{2} \rceil}{\lfloor \frac{4}{2} \rfloor} + \binom{8 - 1 - \lceil \frac{3}{2} \rceil}{\lfloor \frac{3}{2} \rfloor} - (4 - 3) = \binom{6}{2} + \binom{5}{1} - 1 = 19.$$

This means that  $(U_\sigma, \text{orb}(\sigma))$  necessarily belongs to the class **(B)** described in §1.

**Remark 3.3** The upper bound (3.1) will be improved considerably in [5], in the case in which  $P$  is a *simplex*, by using a different combinatorial-topological technique.

## References

- [1] DAIS D. I., HAASE C., ZIEGLER G. M. : *All toric local complete intersection singularities admit projective crepant resolutions*, Tôhoku M. J. **53**, (2001), 95-107.
- [2] DAIS D.I., HAUS U.-U., HENK M.: *On crepant resolutions of 2-parameter series of Gorenstein cyclic quotient singularities*, Results in Math. **33**, (1998), 208-265.
- [3] DAIS D.I., HENK M.: *On a series of Gorenstein cyclic quotient singularities admitting a unique projective crepant resolution*, alg-geom/9803094.
- [4] DAIS D.I., HENK M., ZIEGLER G. M.: *All abelian quotient c.i.-singularities admit projective crepant resolutions in all dimensions*, Adv. in Math. **139**, (1998), 192-239.
- [5] DAIS D.I., HENK M., ZIEGLER G. M.: *On the existence of crepant resolutions of Gorenstein abelian quotient singularities in dimensions  $\geq 4$* , in preparation.
- [6] KALAI G.: *Ridigity and the lower upper bound theorem*, Inventiones Math. **88**, (1987), 125-151.
- [7] ODA T.: *Convex Bodies and Algebraic Geometry*, Springer-Verlag, (1988).
- [8] SCHENZEL P.: *On the number of faces of simplicial complexes and the purity of Frobenius*, Math. Z. **178**, (1981), 125-142.
- [9] STANLEY R.P.: *The upper bound conjecture and Cohen-Macaulay rings*, Studies in Applied Math. **54**, (1975), 135-142.
- [10] STANLEY R.P.: *Hilbert functions of graded algebras*, Advances in Math. **28**, (1978), 57-81.
- [11] STANLEY R.P.: *Combinatorics and Commutative Algebra*, Progress in Math., Vol. **41**, Birkhäuser, (1983); second edition, (1996).
- [12] STANLEY R.P.: *Monotonicity property of  $h$ - and  $h^*$ -vectors*, European Jour. of Combinatorics **14**, (1993), 251-258.
- [13] ZIEGLER G.M.: *Lectures on Polytopes*, GTM, Vol. **152**, Springer-Verlag, (1995).